arXiv:math/0408308v2 [math.QA] 11 Oct 2004

SELBERG INTEGRALS

ALEXANDER VARCHENKO [*],[1]

Abstract. The paper is written for Kluwer's Encyclopaedia of Mathematics.

[*] *Department of Mathematics, University of North Carolina at Chapel Hill,*
Chapel Hill, NC 27599-3250, USA

August, 2004

In 1944 A. Selberg proved the formula [S] :

$$(1)\qquad \int_{\Delta^k[0,1]} \prod_{a=1}^{k} t_a^{\alpha-1}(1-t_a)^{\beta-1} \prod_{1\le a<b\le k}(t_a-t_b)^{2\gamma}\, dt_1\dots dt_k = \prod_{j=0}^{k-1} \frac{\Gamma((j+1)\gamma)}{\Gamma(\gamma)}\, \frac{\Gamma(\alpha+j\gamma)\,\Gamma(\beta+j\gamma)}{\Gamma\Big(\alpha+\beta+(2k-2-j)\gamma\Big)}\ ,$$

where $\Delta^k[x,y] = \{t \in \mathbb{R}^k \mid x \le t_k \le \dots \le t_1 \le y\}$, Re $\alpha > 0$, Re $\beta > 0$, Re $\gamma > \min\{1/k, (\mathrm{Re}\,\alpha)/(k-1), (\mathrm{Re}\,\alpha)/(k-1)\}$. This is a generalization of Euler's formula,

$$\int_0^1 t^{\alpha-1}(1-t)^{\beta-1}\ dt\ =\ \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}\ .$$

The Selberg integral is one of most remarkable multi-dimensional hypergeometric functions with many applications, see references.

Taking a suitable limit of the integral one gets the exponential Selberg integral [M]:

$$\int_{\Delta^k[0,\infty]} \prod_{a=1}^{k} e^{-t_a} t_a^{\alpha-1} \prod_{1\le a<b\le k}(t_a-t_b)^{2\gamma} dt_1\dots dt_k\ =\ \prod_{j=0}^{k-1} \frac{\Gamma((j+1)\gamma)\,\Gamma(\alpha+j\gamma)}{\Gamma(\gamma)}\ .$$

There are the Mellin-Barnes type Selberg integrals [G, TV1]:

$$\int_{-i\infty}^{i\infty}\!\cdots\!\int_{-i\infty}^{i\infty} \prod_{a=1}^{k}\Big(\Gamma(\alpha+t_a)\Gamma(\beta+t_a)\Gamma(\gamma-t_a)\Gamma(\delta-t_a)\ \times$$
$$\prod_{b=1}^{a-1} \frac{\Gamma(t_b-t_a+\epsilon)\Gamma(t_a-t_b+\epsilon)}{\Gamma(t_b-t_a)\Gamma(t_a-t_b)}\Big)\, dt_1\dots dt_k =$$
$$= (2\pi i)^k\ k!\ \prod_{j=0}^{k-1}\Big(\frac{\Gamma((j+1)\epsilon)}{\Gamma(\epsilon)}\ \frac{\Gamma(\alpha+\gamma+j\epsilon)\Gamma(\alpha+\delta+j\epsilon)\Gamma(\beta+\gamma+j\epsilon)\Gamma(\beta+\delta+j\epsilon)}{\Gamma(\alpha+\beta+\gamma+\delta+(2k-2-j)\epsilon)}\Big)\ ,$$

---

[1] Supported in part by NSF grant DMS-0244579.

$$\int_{-i\infty}^{i\infty}\cdots\int_{-i\infty}^{i\infty}\ \prod_{a=1}^{k}\left(u^{2t_a}\Gamma(\alpha+t_a)\Gamma(\alpha-t_a)\ \prod_{b=1}^{a-1}\ \frac{\Gamma(t_b-t_a+\gamma)\Gamma(t_a-t_b+\gamma)}{\Gamma(t_b-t_a)\Gamma(t_a-t_b)}\right)dt_1\dots dt_k =$$
$$(2\pi i)^k\ k!\ (u+u^{-1})^{-k(2\alpha+(k-1)\gamma)}\ \prod_{j=0}^{k-1}\ \frac{\Gamma((j+1)\gamma)\,\Gamma(2\alpha+j\gamma)}{\Gamma(\gamma)}\ ,$$

where $\mathrm{Re}\,\alpha,\beta,\gamma,\delta,\epsilon,\mathrm{u}>0$. In particular,

$$\int_{-i\infty}^{i\infty}\Gamma(\alpha+t)\Gamma(\beta+t)\Gamma(\gamma-t)\Gamma(\delta-t)\ dt\quad=\ 2\pi i\ \frac{\Gamma(\alpha+\gamma)\Gamma(\alpha+\delta)\Gamma(\beta+\gamma)\Gamma(\beta+\delta)}{\Gamma(\alpha+\beta+\gamma+\delta)},$$

$$\int_{-i\infty}^{i\infty}\Gamma(\alpha+t)\Gamma(\alpha-t)\ u^{2t}\ dt\quad=\ 2\pi i\ \Gamma(2\alpha)\ (u+u^{-1})^{-2\alpha},$$

which are formulae for the classical Barnes integrals [WW].

Let $q\ \in\ \mathbb{C},\ 0<|q|<1,\quad (u)_\infty\ =\ \prod_{n=0}^{\infty}(1\ -\ q^n u),\quad \theta(u)=(u)_\infty(qu^{-1})_\infty(q)_\infty$, $\mathbb{T}^k=\{t\in\mathbb{C}^k\ |\ |t_1|=1,\dots,|t_k|=1\}$. The q-Selberg integrals have the form [TV2]:

$$\int_{\mathbb{T}^k}\prod_{a=1}^{k}\frac{\theta(qt_a/\epsilon)\,\theta(u^{k-1}\gamma\delta\epsilon t_a)}{t_a\,(\gamma t_a)_\infty\,(\delta t_a)_\infty\,(\alpha/t_a)_\infty\,(\beta/t_a)_\infty}\prod_{a=1}^{k}\prod_{b=1,\,b\neq a}^{k}\frac{(t_a/t_b)_\infty}{(ut_a/t_b)_\infty}\,dt_1\dots dt_k\ =$$
$$(2\pi i)^k\,k!\ \prod_{j=0}^{k-1}\frac{(u)_\infty\,(u^{k+j-1}\alpha\beta\gamma\delta)_\infty\,\theta(u^j\gamma\epsilon)\,\theta(u^j\delta\epsilon)}{(u^{j+1})_\infty\,(u^j\alpha\gamma)_\infty\,(u^j\beta\gamma)_\infty\,(u^j\alpha\delta)_\infty\,(u^j\beta\delta)_\infty\,(q)_\infty}$$

where $|\alpha|<1,\ |\beta|<1,\ |\gamma|<1,\ |\delta|<1,\ |u|<1,$ and

$$\int_{\mathbb{T}^k}\prod_{a=1}^{k}\frac{\theta(\epsilon t_a)}{t_a\,(\gamma t_a)_\infty\,(\delta t_a)_\infty}\prod_{a=1}^{k}\prod_{b=1,\,b\neq a}\frac{(t_a/t_b)_\infty}{(ut_a/t_b)_\infty}\,dt_1\dots dt_k\ =$$
$$(2\pi i)^k\,k!\ \prod_{j=0}^{k-1}\frac{(u)_\infty\,(u^j\gamma\delta)_\infty\,(qu^j\gamma/\delta)_\infty}{(u^{j+1})_\infty\,(u^j\gamma\delta)_\infty}$$

where $|\gamma|<1,|\delta|<1,|u|<1$. In particular,

$$\int_C\frac{\theta(qt/\epsilon)\,\theta(\gamma\delta\epsilon t)}{(\gamma t)_\infty\,(\delta t)_\infty\,(\alpha/t)_\infty\,(\beta/t)_\infty}\,\frac{dt}{t}\ =\ 2\pi i\,\frac{(\alpha\beta\gamma\delta)_\infty\,\theta(\gamma\epsilon)\,\theta(\delta\epsilon)}{(q)_\infty\,(\alpha\gamma)_\infty\,(\beta\gamma)_\infty\,\infty\,(\alpha\delta)_\infty\,(\beta\delta)_\infty}\ ,$$

where C is an anti-clockwise oriented contour around the origin $t=0$ separating the sets $\{q^s/\gamma,\ q^s/\delta\ |\ s\in\mathbb{Z}_{\leq 0}\},\ \{q^s\alpha,\ q^s\beta\ |\ s\in\mathbb{Z}_{\geq 0}\},$ and

$$\int_C\frac{\theta(\epsilon t)}{(\gamma t)_\infty\,(\delta/t)_\infty}\,\frac{dt}{t}\ =\ 2\pi i\,\frac{(q\gamma/\epsilon)_\infty(\delta\epsilon)_\infty}{(\gamma\delta)_\infty}\ ,$$

where C is an anti-clockwise oriented contour around the origin $t=0$ separating the sets $\{q^s/\gamma\ |\ s\in\mathbb{Z}_{\leq 0}\},\ \{q^s\delta\ |\ s\in\mathbb{Z}_{\geq 0}\}$, see [GR].

One can calculate the integrals over $\mathbb{T}^k$ by residues and get formulas for the q-Selberg Jackson integrals [As, E, Ka1, Ka2, TV2]. Set

$$A(t_1,\dots,t_k;u)\ =\ \prod_{a=1}^{k}\frac{t_a\,(qt_a/\gamma)_\infty\,(qt_a/\delta)_\infty}{(\alpha t_a)_\infty\,(\beta t_a)_\infty}\prod_{1\leq a<b\leq k}\frac{(1-t_b/t_a)\,(qu^{-1}t_b/t_a)_\infty}{(ut_b/t_a)_\infty}\ .$$

If $|qu^{k-1}|<1$, then

$$\sum_{a=0}^{k}\sum_{r_1,\dots,r_k=0}^{\infty}\ (-1)^a\,u^{\sum_{b=1}^{k}(k-b)(k-b+1)r_b-(k-a-1)(k-a)(1+2\sum_{b=1}^{a}r_b)/2}\prod_{c=0}^{k-a-1}\ \frac{\theta(u^{a+c}\gamma/\delta)}{\theta(u^{a-c}\gamma/\delta)}$$
$$\times\ A(q^{r_1}\gamma,q^{r_1+r_2}u\gamma,\dots,q^{r_1+\dots+r_a}u^{a-1}\gamma,q^{r_{a+1}}\delta,\dots,q^{r_{a+1}+\dots+r_k}u^{k-a-1}\gamma;u)\ =$$
$$\prod_{j=0}^{k-1}\frac{(u)_\infty\,(u^{k+j-1}\alpha\beta\gamma\delta)_\infty\,\delta\,\theta(u^j\gamma/\delta)}{(u^{j+1})_\infty\,(u^j\alpha\gamma)_\infty\,(u^j\beta\gamma)_\infty\,(u^j\alpha\delta)_\infty\,(u^j\beta\delta)_\infty}\ .$$

Set

$$B(t_1,\dots,t_k) \;=\; \prod_{a=1}^{k} \tfrac{(qt_a)_\infty}{(\alpha t_a)_\infty} \prod_{1\le a<b\le k} \tfrac{(1-t_b/t_a)\,(qu^{-1}t_b/t_a)_\infty}{(ut_b/t_a)_\infty} \;.$$

If $|v|<\min\,(1,|u^{k-1}|)$, then

$$\sum_{r_1,\dots,r_k=0}^{\infty} v^{\sum_{a=1}^{k}(k-a+1)r_a}\; u^{-\sum_{a=1}^{k}(a-1)(k-a+1)r_a}\; B(q^{r_1},q^{r_1+r_2}u,\dots,q^{r_1+\dots+r_k}u^{k-1}) \;=\; \prod_{j=0}^{k-1} \tfrac{(u)_\infty\,(u^j\alpha v)_\infty\,(q)_\infty}{(u^{j+1})_\infty\,(u^j\alpha)_\infty\,(u^{-j}v)_\infty} \;.$$

Let

$$\theta_1(t,\tau)=-\sum_{j\in\mathbb{Z}} e^{\pi i(j+\frac12)^2\tau+2\pi i(j+\frac12)(t+\frac12)},\qquad \theta_{\kappa,n}(t,\tau)=\sum_{j\in\mathbb{Z}} e^{2\pi i\kappa(j+\frac{n}{2\kappa})^2\tau+2\pi i\kappa(j+\frac{n}{2\kappa})t}$$

where $\kappa,n\in\mathbb{Z}$, $\kappa\ge 2$. Let

$$\sigma_\lambda(t,\tau)=\tfrac{\theta_1(\lambda-t,\tau)\;\theta_1'(0,\tau)}{\theta_1(\lambda,\tau)\;\theta_1(t,\tau)},\qquad E(t,\tau)=\tfrac{\theta_1(t,\tau)}{\theta_1'(0,\tau)},$$

where $'$ denotes the derivative with respect to the first argument. For $k\in\mathbb{Z}_{>0}$, the elliptic Selberg integral is the integral $I_k(\lambda,\tau)=J_k(\lambda,\tau)+(-1)^{k+1}J_k(-\lambda,\tau)$, where

$$J_k(\lambda,\tau) \;=\; \int_{\Delta^k[0,1]} \theta_{2k+2,k+1}(\lambda+\tfrac{1}{k+1}\sum_{a=1}^{k} t_a,\;\tau)\times$$
$$\prod_{a=1}^{k} E(t_a,\tau)^{-k/(k+1)}\sigma_\lambda(t_a,\tau)\;\prod_{1\le a<b\le k} E(t_a-t_b,\tau)^{1/(k+1)}\,dt_1\dots dt_k \;.$$

Then [FSV1, FSV2]

$$I_k(\lambda,\tau) \;=\; c_k\,B_k(\tfrac12+\tfrac{1}{2k+2},-\tfrac{k}{k+1},\tfrac{1}{2k+2})\;\theta_1(\lambda,\tau)^{k+1}\;,$$

where $c_k=(2\pi)^{k/2}e^{\pi ik/(k+1)}e^{-\pi i(k+2)/4}\prod_{a=1}^{k}(e^{-\pi ia/(k+1)}-1)$ and $B_k(\alpha,\beta,\gamma)$ is the Selberg integral in (1).

All previous versions of the Selberg integral are related to representation theory of the Lie algebra sl_2 and appear as solutions to different versions of the KZ equations and dynamical equations, see [FMTV, TV1, TV2, TV3, TV4, TV5, FSV1, FSV2, V]. The following integral is an example of a Selberg type integral associated with sl_3.

Let k_1,k_2 be non-negative integers, $k_1\ge k_2$. For any non-decreasing map $M:\{1,\dots,k_2\}\to\{1,\dots,k_1\}$ such that $M(b)\le k_1-k_2+b$ for any $b=1,\dots,k_2$, introduce a domain $\Delta_M^{k_1,k_2}[x,y]$ in $\mathbb{R}^{k_1+k_2}$ with coordinates $t_1,\dots,t_{k_1},s_1,\dots,s_{k_2}$ defined by the inequalities:

$$x\le t_{k_1}\le\dots\le t_1\le y\,,\qquad x\le s_{k_2}\le\dots\le s_1\le y\,,$$
$$t_{M(b)}\le s_b\le t_{M(b)-1}\,,\qquad b=1,\dots,k_2\,.$$

Here $t_0=y$. For $\gamma\in\mathbb{C}$, consider the chain $C_\gamma^{k_1,k_2}[x,y]=\sum_M X_{M,\gamma}^{k_1,k_2}\Delta_M^{k_1,k_2}[x,y]$ with coefficients

$$X_{M,\gamma}^{k_1,k_2} \;=\; \prod_{b=1}^{k_2} \tfrac{\sin\,(\pi(k_1-k_2-M(b)+b+1)\gamma)}{\sin\,(\pi(k_1-k_2+b)\gamma)} \;.$$

Then [TV5]

$$\begin{aligned}\int_{C_\gamma^{k_1,k_2}[0,1]} \prod_{a=1}^{k_1} t_a^{\alpha-1}(1-t_a)^{\beta_1-1} \prod_{b=1}^{k_2}(1-s_b)^{\beta_2-1} \prod_{a=1}^{k_1}\prod_{b=1}^{k_2} |t_a-s_b|^{-\gamma} \ \times \\ \prod_{1\le a<b\le k_1}(t_a-t_b)^{2\gamma} \prod_{1\le a<b\le k_2}(s_a-s_b)^{2\gamma}\ dt_1\dots dt_{k_1}\, ds_1\dots ds_{k_2} \ = \\ \prod_{j=0}^{k_1-1} \frac{\Gamma(\alpha+j\gamma)\,\Gamma((j+1)\gamma)}{\Gamma(\gamma)} \prod_{j=0}^{k_1-k_2-1} \frac{\Gamma(\beta_1+j\gamma)}{\Gamma(\alpha+\beta_1+(2k_1-k_2-2-j)\gamma)} \ \times \\ \prod_{j=0}^{k_2-1} \frac{\Gamma(\beta_2+j\gamma)\,\Gamma(\beta_1+\beta_2+(j-1)\gamma)\,\Gamma(1+(j-k_1)\gamma)\,\Gamma((j+1)\gamma)}{\Gamma(\beta_2+1+(2k_2-k_1-2-j)\gamma)\,\Gamma(\alpha+\beta_1+\beta_2+(k_1+k_2-3-j)\gamma)\,\Gamma(\gamma)}\ .\end{aligned}$$

In the k-dimensional Euclidean space $\mathbb{R}^k$ consider the reflection hyperplanes of a Coxeter group G. Let $P(t)$ be the product of distances of the point $t=(t_1,...,t_k)$ from all reflection hyperplanes associated with G. Let N be the number of hyperplanes. Then [Ma, Op, M]

$$\int_{\mathbb{R}^k} e^{-\sum t_i^2/2}\, |P(t)|^{2\gamma}\, dt_1\dots dt_k \ = \ 2^{-N\gamma}\, (2\pi)^{k/2} \prod_{j=1}^{k} \frac{\Gamma(1+\gamma d_j)}{\Gamma(1+\gamma)}\ ,$$

where d_j are the degrees of basis polynomials in the space of homogeneous polynomials, which are invariant with respect to G. The q-analogs of this formula were proved in [Ch].

Here is an example of a Selberg type integral involving the elliptic gamma function, see [Sp]. Let

$$\Gamma(t;p,q) \ = \ \prod_{j,k=0}^{\infty} \frac{1-t^{-1}q^{j+1}p^{k+1}}{1-tq^jp^k}$$

be the elliptic gamma function and $(q;q)_\infty = \prod_{n=1}^\infty (1-q^n)$. For complex parameters $u_0,\dots,u_4$, set $A=\prod_{m=0}^4 u_m$. Assume that $|p|<1, |q|<1, |u_m|<1, |pq|<|A|$. Then

$$\frac{1}{2\pi i}\int_{\mathbb{T}^1} \frac{\prod_{m=0}^4 \Gamma(tu_m;p,q)\,\Gamma(t^{-1}u_m;p,q)}{\Gamma(t^2;p,q)\,\Gamma(t^{-2};p,q)\,\Gamma(tA;p,q)\,\Gamma(t^{-1}A;p,q)}\,\frac{dt}{t} \ = \ \frac{2\prod_{0\le l<m\le 4}\Gamma(u_lu_m;p,q)}{(q;q)_\infty\,(p;p)_\infty \prod_{m=0}^4 \Gamma(Au_m^{-1};p,q)}\ .$$

References

[A1] K. Aomoto, *Jacobi polynomials associated with Selberg integrals*, SIAM J. Math. Anal. **18** (1987), 545–549.

[A2] K. Aomoto, *On the complex Selberg integral*, Q. J. Math. Oxford **38** (1987), 385–399.

[AK] K. Aomoto and Y. Kato, *Gauss decomposition of connection matrices for symmetric A-type Jackson integrals*, Selecta Math., New Series **1** (1995), 623–666.

[As] R. Askey, *Some basic hypergeometric extensions of integrals of Selberg and Andrews*, SIAM J. Math. Anal., **11** (1980), 938–951.

[AAR] G. Andrews, R. Askey, and R. Roy, *Special functions*, Encyclopedia of Mathematics and its Applications, **71**, Cambridge University Press, Cambridge, 1999. xvi+664 pp.

[Ch] I. Cherednik, *Difference Macdonald-Mehta conjecture*, Internat. Math. Res. Notices (1997), No. 10, 449–467.

[DF] V. Dotsenko and V. Fateev *Four-point correlation functions and the operator algebra in 2-D conformal invariant theories with central charge $C \le 1$*, Nucl. Phys. B **251** (1985), 691–734.

[E] R. Evans, *Multidimensional beta and gamma integrals*, Contemporary Mathematics, **166** (1994), 341–357.

[FMTV] G. Felder, Y. Markov, V. Tarasov, and A. Varchenko, *Differential Equations Compatible with KZ Equations*, QA/0001184, Journal of Math. Phys., Analysis and Geometry, 3 (2000), 139–177.

[FSV1] G. Felder, L. Stevens, and A. Varchenko, *Modular transformations of the elliptic hypergeometric functions, Macdonald polynomials, and the shift operator*, Mosc. Math. J. **3** (2003), no. 2, 457–473.

[FSV2] G. Felder, L. Stevens, and A. Varchenko, *Elliptic Selberg integrals and conformal blocks*, Math. Res. Lett. **10** (2003), no. 5-6, 671–684.

[FV] G. Felder and A. Varchenko, *The elliptic gamma function and* $\mathrm{SL}(3,\mathbb{Z}) \ltimes Z^3$, Adv. Math. **156** (2000), no. 1, 44–76.

[GR] G. Gasper and M. Rahman, *Basic hypergeometric series*, Encycl. Math. Appl., Cambridge University Press, 1990.

[G] R.A. Gustafson, *A generalization of Selberg's beta integral*, Bull. Amer. Math. Society **22** (1990), 97–105.

[Ka1] K. Kadell, *A proof of Askey's conjectured q-analogue of Selberg's integral and a conjecture of Morris*, SIAM J. Math. Anal., **19** (1988) 969–986.

[Ka2] K. Kadell, *A simple proof of an Aomoto-type extension of Askey's last conjectured Selberg q-integral*, J. Math. Anal. Appl., **261** (2001), no. 2, 419–440.

[M] M.L. Mehta, *Random Matrices*, Academic Press, 1991.

[Ma] I. Macdonald, *Some conjectures for root systems*, SIAM J. Math. Anal., 13:6 (1982), 988–1007.

[N] Y. Neretin, *Rayleigh triangles and non-matrix interpolation of matrix beta-integrals*, (Russian) Mat. Sb. **194** (2003), no. 4, 49–74; translation in Sb. Math. **194** (2003), no. 3-4, 515–540.

[Op] E. Opdam, *Some applications of hypergeometric shift operators*, Invent. Math. **98** (1989), no. 1, 1–18.

[S] A. Selberg, *Bemerkninger om et multiplet integral*, Norsk Mat. Tidsskr. **26** (1944), 71–78.

[Sp] V. Spiridonov, *An elliptic beta integral*, New trends in difference equations (Temuco, 2000), 273–282, Taylor and Francis, London, 2002.

[TV1] V. Tarasov and A. Varchenko, *Geometry of q-hypergeometric functions as a bridge between Yangians and quantum affine algebras*, Invent. Math. **128** (1997), no. 3, 501–588.

[TV2] V. Tarasov and A. Varchenko, *Geometry of q-hypergeometric functions, quantum affine algebras and elliptic quantum groups*, Astrisque No. **246** (1997), vi+135.

[TV3] V. Tarasov and A. Varchenko, *Difference equations compatible with trigonometric KZ differential equations*, Internat. Math. Res. Notices 2000, no. 15, 801–829.

[TV4] V. Tarasov and A. Varchenko, *Duality for Knizhnik-Zamolodchikov and dynamical equations*, The 2000 Twente Conference on Lie Groups (Enschede), Acta Appl. Math. **73** (2002), no. 1-2, 141–154.

[TV5] V. Tarasov and A. Varchenko, *Selberg-type integrals associated with* sl_3, Lett. Math. Phys. **65** (2003), no. 3, 173–185.

[V] A. Varchenko, *Special functions, KZ type equations, and representation theory*, CBMS Regional Conference Series in Mathematics, **98**, AMS, 2003.

[WW] E.T. Whittaker and G.N. Watson, *A Course of Modern Analysis*, Cambridge University Press, 1927.